



# Epidemiological Trends of HIV/AIDS in Sub-Saharan Africa: A 2024 Perspective

Muhindo Anitah

Faculty of Pharmacy Kampala International University Uganda

Email: [anitah.muhindo@studwc.kiu.ac.ug](mailto:anitah.muhindo@studwc.kiu.ac.ug)

## ABSTRACT

The HIV/AIDS epidemic in Sub-Saharan Africa remains a major public health challenge, with the region accounting for approximately 68% of the global population living with HIV. This review explores the current epidemiological trends in 2024, highlighting the demographic shifts, socioeconomic factors, and public health interventions shaping the epidemic. Despite progress in reducing new infections and expanding antiretroviral therapy (ART) coverage, significant challenges remain, particularly among vulnerable populations such as women, adolescents, and key populations like men who have sex with men (MSM) and sex workers. The review also discusses the impact of public health interventions, such as the prevention of mother-to-child transmission (PMTCT) and the integration of HIV services with other health programs. Socioeconomic barriers, stigma, and disparities in healthcare access are critical issues that must be addressed to achieve long-term control of the epidemic. The evolving landscape of HIV/AIDS, including emerging issues like aging populations and co-infection with tuberculosis, requires innovative approaches to prevention and treatment.

**Keywords:** HIV/AIDS, Sub-Saharan Africa, epidemiological trends, antiretroviral therapy (ART), mother-to-child transmission (MTCT),

## INTRODUCTION

The HIV/AIDS epidemic in Sub-Saharan Africa is a significant burden, with approximately 68% of the world's total population living with HIV (PLHIV) residing in this region. This underscores the pressing nature of the epidemic, which poses considerable health, social, and economic challenges across various countries. The impact of HIV/AIDS in Sub-Saharan Africa is multifaceted, influencing not only the health of individuals but also the broader socio-economic fabric of communities [1]. Health challenges include high prevalence rates, leading to significant morbidity and mortality, particularly among vulnerable populations. The World Health Organization estimates that over 1.3 million people died from AIDS-related illnesses in the region in 2021 [2]. The ongoing epidemic strains healthcare systems, many of which are already overburdened and under-resourced, complicating the delivery of essential services beyond HIV care [3]. The epidemic also intersects with other health issues, notably tuberculosis (TB), which remains a leading cause of death among PLHIV. Coinfection exacerbates health outcomes and complicates treatment protocols, creating a pressing need for integrated care approaches that can simultaneously address both conditions.

Social challenges include stigma and discrimination against PLHIV, leading to social isolation and limiting individuals' access to healthcare, education, and employment opportunities. This stigma is often rooted in cultural beliefs and misinformation about the transmission and nature of the virus, which can deter individuals from seeking testing and treatment, ultimately perpetuating the cycle of infection [4]. The epidemic also disproportionately affects specific demographic groups, including women, adolescents, and key populations such as

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

men who have sex with men (MSM) and sex workers. Economic challenges include reduced productivity, increased healthcare costs, and loss of human capital. Families affected by HIV often face financial hardship due to medical expenses and loss of income from sick family members [5]. In many cases, this leads to a cycle of poverty, where the economic burden of the disease inhibits access to education and healthcare, further entrenching communities in socio-economic challenges. Public health interventions have been implemented, with varying degrees of success. The introduction of antiretroviral therapy (ART) has revolutionized HIV management, significantly reducing morbidity and mortality among PLHIV [6]. However, gaps remain in access to these services, particularly among marginalized populations. Ensuring that prevention and treatment services reach all segments of the population is critical for achieving epidemic control. Emerging issues are influencing the landscape of HIV/AIDS in Sub-Saharan Africa, such as the COVID-19 pandemic, mental health, and research on HIV transmission dynamics. Future efforts must prioritize equity, innovation, and sustainability to build resilient health systems capable of addressing the ongoing challenges of HIV/AIDS in the region [7].

#### Current Prevalence Rates

As of 2024, Sub-Saharan Africa is the epicenter of the global HIV/AIDS epidemic, with an estimated 26 million people living with HIV (PLHIV) in the region. This accounts for approximately 68% of the global total. The adult prevalence rate stands at around 5%, indicating a significant proportion of the population is affected by the virus [8]. Despite ongoing efforts in prevention and treatment, the burden of HIV/AIDS remains substantial, with some countries experiencing particularly high rates of infection. Regional disparities exist within Sub-Saharan Africa, with South Africa having the highest number of PLHIV in the world at about 19.1%. Nigeria, the most populous country in Africa, also has one of the largest HIV populations, with an adult prevalence rate of about 1.5%. Kenya reports an adult prevalence rate of around 4.9%, with significant regional variation [9]. The country has made progress in reducing new infections but still faces challenges related to stigma, access to healthcare, and social determinants of health.

While there have been encouraging declines in new HIV infections in several countries due to enhanced prevention efforts, the overall burden of the epidemic remains high, particularly in eastern and southern Africa. Countries like Eswatini and Botswana have reported substantial reductions in new infections due to successful public health interventions, including widespread testing, antiretroviral therapy (ART), and community-based education initiatives. However, eastern and southern Africa continue to bear the brunt of the epidemic, necessitating sustained prevention and treatment efforts to avoid resurgence [10]. Key populations within Sub-Saharan Africa remain disproportionately affected by HIV, contributing to the overall prevalence rates. These groups include men who have sex with men (MSM), sex workers, people who inject drugs, and transgender individuals. Stigma, discrimination, and legal barriers often hinder access to testing and treatment for these populations. Women and adolescents face higher risks of infection due to biological vulnerability and socio-economic factors. The current landscape of HIV prevalence in Sub-Saharan Africa as of 2024 reflects a region still grappling with the complexities of the epidemic. Addressing the diverse needs of key populations and strengthening health systems will be crucial to mitigating the impact of HIV/AIDS and achieving long-term control of the epidemic [11].

#### Demographic Shifts

Recent studies have shown significant demographic shifts in the epidemiology of HIV/AIDS across Sub-Saharan Africa, highlighting the evolving nature of the epidemic. Traditional risk groups such as men who have sex with men (MSM), sex workers, and people who inject drugs (PWID) continue to be disproportionately affected, but emerging trends indicate that the virus is increasingly impacting older adults and women. These demographic changes necessitate a reevaluation of public health strategies and interventions to address the specific needs of diverse population groups. Key populations include MSM, which remain one of the most vulnerable in the context of HIV/AIDS due to stigma, discrimination, and criminalization in many Sub-Saharan African countries [12]. Tailored interventions, including targeted outreach programs, peer education, and safe-sex practices, are essential to reduce transmission within this demographic. Female sex workers are another key population disproportionately affected by HIV, and interventions aimed at this group have been effective in some regions, particularly those that integrate HIV prevention with broader sexual and reproductive health services. Empowering sex workers through legal protection, healthcare access, and community support is crucial in mitigating their risk of infection.

People who inject drugs (PWID) face unique challenges related to HIV transmission, primarily due to needle-sharing practices. Harm reduction strategies, including needle exchange programs and opioid substitution therapy, have shown promise in reducing new infections among this demographic [13]. However, the stigma surrounding

**This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**

drug use often limits access to these essential services, underscoring the need for comprehensive policy reforms to support PWID. The impact on older adults is noticeable, with an increase in infections among those aged 50 and above. This shift can be attributed to factors such as increased longevity of PLHIV, risk behaviors, limited awareness, and gender inequality. Women, particularly those of reproductive age, continue to bear a significant burden of HIV, and efforts to reduce mother-to-child transmission (MTCT) through antenatal care and access to ART during pregnancy and breastfeeding have shown effectiveness but still require extensive reach to ensure all women receive necessary care [14]. Emerging demographics include adolescents and young people, who represent a significant proportion of new infections. Issues such as lack of comprehensive sexual education, social stigmas, and intersectionality must be addressed to develop effective, inclusive interventions.

#### **Socioeconomic Factors and Risk**

Socioeconomic factors significantly impact the HIV/AIDS epidemic in Sub-Saharan Africa, affecting the risk of transmission and the ability of individuals and communities to access effective prevention and treatment services. Factors such as poverty, lack of education, limited access to healthcare, and social stigmatization contribute to increased vulnerability to HIV and barriers to testing, treatment, and long-term care. Poverty and economic disparities increase susceptibility to HIV by restricting access to healthcare facilities, increasing risk behaviors, and limiting treatment affordability. Lack of education and awareness also contribute to the spread of HIV, with many people lacking basic knowledge about HIV transmission and prevention methods. This lack of knowledge affects young people, leading to higher rates of new infections and risky behaviors [15]. Gender disparities in education further limit women's ability to make informed decisions about their sexual and reproductive health. Access to healthcare services is a crucial factor in HIV outcomes in Sub-Saharan Africa. The region faces challenges such as underfunded health systems, shortages of healthcare workers, and inadequate infrastructure. Geographic barriers, healthcare system capacity, and disparities in urban vs. rural access contribute to poor health outcomes.

Social stigma and discrimination against people living with HIV/AIDS (PLHIV) also hinder effective prevention, treatment, and care. Fear of disclosure, healthcare discrimination, and mental health implications further exacerbate health disparities. High levels of mobility and migration within and between countries contribute to the spread of HIV, complicating prevention and control efforts. Labor migration, cross-border spread, and challenges in continuity of care are significant challenges for mobile populations. Gender inequality and social norms exacerbate the risk of HIV infection, particularly for women and girls. Power imbalances, early marriage and childbearing, and maternal HIV transmission further complicate the burden of HIV. Socioeconomic factors such as poverty, lack of education, limited healthcare access, stigma, and high levels of mobility play a critical role in shaping the epidemiology of HIV/AIDS in Sub-Saharan Africa [16]. Addressing these factors through comprehensive public health strategies is essential for reducing HIV transmission and improving the quality of life for PLHIV. Interventions must be tailored to the unique challenges posed by different socioeconomic contexts, with a particular focus on expanding healthcare access, promoting gender equality, and reducing stigma and discrimination across all levels of society.

#### **Impact of Public Health Interventions**

Public health interventions have significantly impacted the HIV/AIDS epidemic in Sub-Saharan Africa, with the introduction of antiretroviral therapy (ART) and targeted prevention programs transforming individual and community-level responses. However, significant gaps remain, particularly regarding access to treatment, adherence, and effective intervention strategies in rural and underserved populations [17]. Antiretroviral therapy (ART) has been a game changer for PLHIV, leading to improved survival rates and reduced transmission. However, challenges in access persist, particularly in rural and remote areas due to inadequate healthcare infrastructure, shortages of trained healthcare workers, limited access to testing facilities, and inconsistent drug supplies. Treatment adherence remains a challenge, and community-based support systems and counseling services are essential for maintaining ART effectiveness.

Prevention programs have evolved alongside treatment efforts, focusing on high-risk populations and key drivers of the epidemic. New methods have been introduced to address specific challenges and target vulnerable populations, such as condom distribution and promotion, pre-exposure prophylaxis (PrEP), harm reduction for PWID, and prevention of mother-to-child transmission (PMTCT). Despite these successes, gaps remain in scaling PMTCT programs to all parts of Sub-Saharan Africa, as many women do not have access to antenatal care and a lack of infrastructure can make it difficult to reach pregnant women with necessary interventions. Addressing gaps in rural and underserved areas is crucial, as they often lack the basic healthcare infrastructure necessary to support HIV testing, treatment, and prevention services [18]. Healthcare workforce shortages, stigma, and discrimination

**This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**

also pose barriers to accessing necessary services. The HIV epidemic in Sub-Saharan Africa is evolving, presenting new public health challenges and opportunities. Interventions must address adolescent and youth populations, addressing risk factors like early sexual initiation and limited access to sexual health education. The intersection of HIV and non-communicable diseases is becoming an emerging public health concern, and healthcare systems must manage aging PLHIV patients. Integrating HIV and TB services is crucial for reducing TB-related mortality. Future public health strategies should focus on expanding access to ART, improving treatment adherence, addressing stigma, and targeting emerging issues.

#### **Trends in Mother-to-Child Transmission**

Mother-to-child transmission (MTCT) of HIV remains a significant public health issue in Sub-Saharan Africa, with approximately 160,000 new pediatric HIV infections occurring annually. MTCT can occur during pregnancy, childbirth, or breastfeeding, and without intervention, the transmission rate can be as high as 30-45%. However, with timely interventions, particularly the use of antiretroviral therapy (ART), transmission rates can be reduced to below 5%. Over the past two decades, significant progress has been made in PMTCT programs across Sub-Saharan Africa, particularly in countries with high HIV prevalence [8]. Key achievements include increased ART coverage, a decline in pediatric infections, and improved maternal health. However, challenges such as late presentation to antenatal care, insufficient ART coverage, breastfeeding transmission, and HIV testing and linkage to care continue to hinder the full elimination of MTCT in Sub-Saharan Africa. Emerging trends and issues in MTCT include a focus on retention in care, which aims to ensure that women are enrolled in PMTCT programs after delivery. Postnatal follow-up and ART adherence are crucial for preventing breastfeeding-related transmission of HIV in Sub-Saharan Africa. Many countries are integrating PMTCT services with maternal, newborn, and child health (MNCH) services to improve care outcomes. Adolescent girls and young women are disproportionately affected by HIV, and targeted interventions for them are essential to reduce MTCT rates [11]. Innovative approaches to ART delivery, such as community-based ART distribution and mobile health technologies, are being piloted to improve access and adherence to treatment. To further reduce MTCT rates and eliminate pediatric HIV in Sub-Saharan Africa, early HIV diagnosis, improved ART coverage and retention, and addressing social and structural barriers are essential. Community-based interventions can reduce stigma and encourage supportive environments for women accessing PMTCT services. Expanding PMTCT programs beyond pregnancy and delivery to include comprehensive postnatal care is also crucial for preventing transmission during breastfeeding. By strengthening PMTCT programs and expanding innovative approaches, Sub-Saharan Africa can move closer to achieving the elimination of pediatric HIV.

#### **Co-Infections and Emerging Challenges**

Co-infections with communicable diseases, particularly tuberculosis (TB), pose significant challenges in HIV/AIDS management in Sub-Saharan Africa. The region faces a dual burden of both HIV and TB due to weakened immune systems of individuals living with HIV. This HIV-TB syndemic exacerbates health outcomes and increases morbidity and mortality. The rise of drug-resistant strains of HIV and TB has emerged as a critical challenge, complicating treatment regimens and emphasizing the need for robust surveillance and research [9]. Tuberculosis is the most common co-infection among people living with HIV (PLHIV) in Sub-Saharan Africa, and it is the leading cause of death among PLHIV worldwide. The HIV-TB co-infection dynamic creates several pressing public health challenges, including increased mortality, diagnostic challenges, treatment complications, and immune reconstitution inflammatory syndrome (IRIS). Drug-resistant TB (DR-TB) is another major challenge in the management of co-infections in Sub-Saharan Africa. Multidrug-resistant TB (MDR-TB) and extensively drug-resistant TB (XDR-TB) are particularly concerning, as they are more difficult and expensive to treat and often have poorer outcomes. The emergence of drug-resistant HIV poses another significant challenge in the HIV epidemic in Sub-Saharan Africa [16]. The increased prevalence of drug-resistant HIV strains, particularly among individuals who have interrupted treatment or failed to adhere to ART regimens, makes it more difficult to select effective first-line treatments. Co-infections, particularly with tuberculosis, and the emergence of drug-resistant strains of HIV and TB present major challenges to the effective management of HIV/AIDS in Sub-Saharan Africa. Effective management requires ongoing surveillance and drug resistance testing, which are not always available in resource-constrained areas. Strengthening surveillance systems and ensuring healthcare providers have the tools and resources to detect and manage drug resistance is crucial to prevent further transmission of resistant strains. In addition to TB, PLHIV in Sub-Saharan Africa are at risk of co-infections with other opportunistic infections, such as malaria, hepatitis B and C, and human papillomavirus (HPV). Integrated care approaches are essential to improve health outcomes for PLHIV, including one-stop clinics, community-based care, and strengthening surveillance and research. Expanding access to second-line

**This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**

treatments, improving diagnostic capacity, and investing in community-based care are critical strategies for reducing morbidity and mortality in PLHIV and curbing the spread of co-infections.

### CONCLUSION

The epidemiological trends of HIV/AIDS in Sub-Saharan Africa, as of 2024, reflect both progress and ongoing challenges in the fight against the epidemic. While significant strides have been made in reducing new infections, particularly through widespread antiretroviral therapy (ART) and targeted prevention programs, the region continues to bear a disproportionate burden of the global epidemic. Socioeconomic factors, such as poverty, limited healthcare access, and persistent stigma, exacerbate the situation, particularly among key populations like men who have sex with men (MSM), sex workers, people who inject drugs (PWID), women, and adolescents.

Demographic shifts, including rising infections among older adults and women, highlight the need for tailored public health strategies that address the diverse needs of the population. Socioeconomic disparities, gender inequality, and healthcare access gaps continue to influence the epidemic's trajectory, underscoring the importance of strengthening healthcare systems, expanding education, and reducing stigma. Mother-to-child transmission (MTCT) rates have seen substantial declines thanks to effective prevention programs, yet gaps remain, particularly in rural and underserved areas. Integrating PMTCT services with broader maternal and child health initiatives and improving ART coverage and retention are critical for further reducing pediatric infections. Moving forward, future strategies must focus on expanding access to ART, addressing treatment adherence challenges, and ensuring equitable healthcare services for all affected populations. The intersection of HIV with other health issues like tuberculosis and non-communicable diseases also requires integrated care approaches. To control and ultimately end the HIV/AIDS epidemic in Sub-Saharan Africa, a continued focus on innovation, equity, and sustainability will be essential.

### REFERENCES

1. Alum, E. U., Obeagu, E. I., Ugwu, O. P. C., Samson, A. O., Adepoju, A. O., Amusa, M. O. Inclusion of nutritional counseling and mental health services in HIV/AIDS management: A paradigm shift. *Medicine (Baltimore)*. 2023;102 (41):e35673. <http://dx.doi.org/10.1097/MD.00000000000035673>. PMID: 37832059; PMCID: PMC10578718.
2. Obeagu, E.I., Alum, E.U. and Obeagu, G.U. Factors Associated with Prevalence of HIV Among Youths: A Review of Africa Perspective. *Madonna University Journal of Medicine and Health Sciences*, 2023;3(1): 13-18. <https://madonnauniversity.edu.ng/journals/index.php/medicine>
3. UNAIDS. (2023). *Global HIV & AIDS Statistics — 2023 Fact Sheet*. UNAIDS. Retrieved from <https://www.unaids.org>
4. World Health Organization (WHO). (2022). *HIV/AIDS in Sub-Saharan Africa: Progress towards 2030 Targets*. World Health Organization Regional Office for Africa. Available at <https://www.afro.who.int>
5. Alum, E. U., Uti, D. E., Ugwu, O. P., Alum, B. N. Toward a cure - Advancing HIV/AIDS treatment modalities beyond antiretroviral therapy: A Review. *Medicine (Baltimore)*. 2024 Jul 5;103(27):e38768. doi: 10.1097/MD.00000000000038768. PMID: 38968496.
6. Kharsany, A. B. M., & Karim, Q. A. (2022). HIV Infection and AIDS in Sub-Saharan Africa: Current Status, Challenges, and Opportunities. *The Open AIDS Journal*, 16, 1-10. DOI: 10.2174/1874613602216010001
7. Alum, E. U., Ugwu, O. P. C., Obeagu, E. I., Aja, P. M., Okon, M. B., Uti, D. E. Reducing HIV Infection Rate in Women: A Catalyst to reducing HIV Infection pervasiveness in Africa. *International Journal of Innovative and Applied Research*. 2023; 11(10):01-06. DOI:10.58538/IJIAR/2048. <http://dx.doi.org/10.58538/IJIAR/2048>
8. Obeagu, E. I., Nwosu, D. C., Ugwu, O. P. C. and Alum, E. U. Adverse Drug Reactions in HIV/AIDS Patients on Highly Active Antiretro Viral Therapy: A Review of Prevalence. *NEWPORT INTERNATIONAL JOURNAL OF SCIENTIFIC AND EXPERIMENTAL SCIENCES (NIJSES)*. 2023; 4(1):43-47. <https://doi.org/10.59298/NIJSES/2023/10.6.1000>
9. Mills, E. J., & Beyrer, C. (2022). The Future of HIV in Africa: Sustainable Prevention and Treatment Solutions. *Lancet HIV*, 9(5), e301-e308. DOI: 10.1016/S2352-3018(22)00039-3
10. Gupta, G. R., et al. (2023). Global Inequalities in HIV, TB, and Malaria: Addressing Disparities to Achieve Health Equity. *Journal of Global Health*, 13, 07005. DOI: 10.7189/jogh.13.07005
11. UNAIDS & WHO. (2024). *2024 Epidemiological Update on HIV in Sub-Saharan Africa: Emerging Trends and Challenges*. Joint United Nations Programme on HIV/AIDS (UNAIDS) & World Health Organization (WHO).

12. Bekker, L. G., Venter, F., & Myer, L. (2023). Changing the Narrative: Epidemiology and Response to the HIV Epidemic in South Africa. *PLOS Medicine*, 20(6), e1004141. DOI: 10.1371/journal.pmed.1004141
13. Alum, E. U., Obeagu, E. I., Ugwu, O. P.C., Aja, P. M. and Okon, M. B. HIV Infection and Cardiovascular diseases: The obnoxious Duos. *Newport International Journal of Research in Medical Sciences (NIJRMS)*, 2023; 3(2): 95-99. <https://nijournals.org/wp-content/uploads/2023/07/NIJRMS-3-295-99-2023.pdf>.
14. Alum, E. U., Ugwu, O. P.C., Obeagu, E. I. and Okon, M. B. Curtailing HIV/AIDS Spread: Impact of Religious Leaders. *Newport International Journal of Research in Medical Sciences (NIJRMS)*, 2023; 3(2): 28-31. <https://nijournals.org/wp-content/uploads/2023/06/NIJRMS-32-28-31-2023-rm.pdf>
15. Avert. (2023). HIV and AIDS in East and Southern Africa: Regional Overview. Avert. Available at <https://www.avert.org>
16. Heidari, S., et al. (2023). Impact of COVID-19 on HIV Prevention, Testing, and Treatment in Sub-Saharan Africa. *AIDS Reviews*, 25(1), 50-62. DOI: 10.24875/AIDSRev.M22000050
17. Alum, E. U., Obeagu E. I., Ugwu, O, P.C., Egba S, I., Uti, D, E., Ukaidi, C, U, A., Echegu, D, A., Confronting Dual Challenges: Substance Abuse and HIV/AIDS. *Elite Journal of HIV*, 2024; 2(5): 1-8. <https://epjournals.com/journals/EJHIV>
18. Eshleman, S. H., & Serwadda, D. (2022). Long-Term ART Outcomes in Sub-Saharan Africa: New Insights from Population-Based Cohorts. *Journal of Acquired Immune Deficiency Syndromes (JAIDS)*, 90(4), 505-515. DOI: 10.1097/QAI.0000000000003033

**CITE AS: Muhindo Anitah (2024). Epidemiological Trends of HIV/AIDS in Sub-Saharan Africa: A 2024 Perspective. *Research Output Journal of Public Health and Medicine* 4(3):1-6. <https://doi.org/10.59298/ROJPHM/2024/431600>**